

Colorado Coalition for School IPM Newsletter

December 2018

Inside this issue:

- Partner Spotlight: North Central IPM Center
- School Spotlight:
- Featured Pest of the Month: Indian Meal Moth
- Current Pests

Colorado Coalition for School IPM Agency Partner Spotlight: North Central IPM Center

Stop School Pests IPM Training Renamed The Pest Defense for Healthy Schools

By

North American IPM Institute and North Central IPM Center

The [National School IPM Working Group](#)'s flagship free online training, Stop School Pests, has been renamed The Pest Defense for Healthy Schools to emphasize its focus on creating healthy, safe spaces for students and school staff. The rebranding is based on outcomes from a national survey of nearly 100 individuals working in schools.

Pests can damage infrastructure, spread disease and produce asthma-triggering allergens. Due to their behavior and biology, school-aged children are particularly susceptible to exposure to both pests and pesticides.

[The Pest Defense](#) is free and includes individualized educational modules for custodians, maintenance, facility managers, food service professionals, teachers, administrators, nurses and grounds professionals. Modules address how these professionals and others can help prevent pest problems as they go about their

daily activities. For example, just ensuring exterior doors have sweeps that effectively close the gap between the bottom of the door and the door sill can reduce pest complaints by up to 65%!

Modules address tactics for preventing and eliminating pests that can be found in school buildings or on school grounds including head lice, bed bugs, rodents, weeds and more. The training was developed with support from [US EPA](#) and the [North Central IPM Center](#).

To learn more about how your school can implement this great resource or to sponsor a local training, please contact [Julian Cooper](#).

Read the official press release [here](#).

CCSIPM Spotlight: 2018 President's Environmental Youth Award (PEYA)

EPA Announces 2018 President's Environmental Youth Award

Applications are now being accepted for the 2018 President's Environmental Youth Award (PEYA).



The PEYA program recognizes outstanding environmental stewardship projects by K-12 youth, promoting awareness of our nation's natural resources and encouraging positive community involvement.

The PEYA program celebrates student leadership in service projects to protect the environment and keep our global community healthy. Winners of this year's awards will be invited to a ceremony in Washington, D.C. in mid-2019, and have their project mentioned on EPA's website.

EPA will select up to two winners in each of EPA’s 10 Regions – one regional winner for Grades K-5 and one regional winner for Grades 6-12.

All student projects must be sponsored by at least one adult over the age of 21. The application and eligibility information are available at: <https://www.epa.gov/education/presidents-environmental-youth-award>.

Applications are due February 1, 2019.

PEYA is an annual award administered by the Office of Environmental Education at EPA. Since 1971, EPA has recognized young people for protecting our nation’s air, water, land, and ecology. Each year the PEYA program honors a wide variety of projects developed by young individuals, school classes (kindergarten through high school), summer camps, public interest groups, and youth organizations to promote environmental awareness. Through environmental education and stewardship activities, students develop the critical thinking skills experience to make informed decisions and take responsible actions to address difficult environmental issues.

For more information, please contact PEYA@epa.gov.

For information on recent winners, visit: <https://www.epa.gov/newsreleases/epa-honors-award-winning-environmental-education-teachers-and-students>

Featured Pest of the Month: Indian Meal Moth

The Indian Meal Moth

The Indian meal moth (*Plodia interpunctella*) is a small, grayish-brown the most common household moth that can reproduce in Colorado homes. It develops as a pest of various foods commonly found in pantries. The caterpillars can seriously damage susceptible food items and the adult moths can become annoying as they fly through the home.

All stages of the Indian meal moth may be found in homes. The adult is a small moth, about 3/8 inch long with a wing span of about 5/8 inch. The overall body color is generally dirty gray but the tip half of the wing is rusty



Figure 1. Adult Indian meal moth. (Photo from the K. Gray collection.)



Figure 2: Indian meal moths cause problems on dried food products in the home. Adult and larvae shown.

brown or nearly bronze. This wing marking pattern allows Indian meal moth to be easily distinguished from other household moths.

The caterpillar stage is usually cream colored, sometimes with yellowish-green or pinkish shades, and has a dark brown head. Normally they stay associated with foods, but the full grown caterpillars, about 2/3 inch long, may be seen as they wander in search of a place to pupate.

Life History and Habits

Indian meal moths develop in many kinds of stored foods. Coarsely ground grains and cereal products are commonly infested. Dried herbs, dried fruits, and nuts are also highly favored. Pet foods such as dried dog food, flaked fish food, and bird seed can also become infested. Indian meal moth may also breed in ornamental items made of dried flowers or seeds.



Figure 3. Full-grown larva and pupa of Indian meal moth. (Photo by W. Cranshaw.)

The moths usually fly at dusk and through the night. Females lay tiny eggs (ca. 0.5 mm) on or near potential food items. The newly hatched caterpillars (larvae) seek out foods and begin to develop. As they feed they often produce silk that loosely binds to food fragments. In large, undisturbed containers feeding is



Figure 4. Indian meal moth larva. (Photo from the K. Gray collection.)

concentrated on the surface as the larvae do little burrowing. However, caterpillars may occur throughout the product within small, loose packages typically found in household pantries. The caterpillars are capable of chewing through plastic bags and thin cardboard.

The rate of development is dependent on factors including temperature and food quality. Under optimum temperatures the caterpillars may become fully-grown in as little as a month; development typically requires a much longer period. When feeding is completed the caterpillars usually wander from the food and search for a place to pupate. They

then create a loose cocoon covering within which they pupate and subsequently transform to the adult stage. Cocoons are most often located in cracks or confined spaces, such as the junction between walls and ceiling.

The adults emerge, mate and lay eggs. Since the moths do not feed they usually survive little more than a week. During this time the female may deposit over 200 eggs, laid either directly on food or in crevices adjacent to the stored foods on which they feed.

Probably three or four generations can be completed annually in a Colorado home although all stages may be present as generations overlap. For reasons that are unclear, adults are usually observed most commonly from November through February.

Management of Indian Meal Moth

The first, and most critical, step is to identify all sources of infestation. The presence of some webbing is usually the most effective way to determine which items are infested.

This examination must be thorough as the range of materials potentially infested is so broad. First go through items in the pantry which may host Indian meal moth. They are commonly found in coarse cereal products (e.g. oatmeal, breakfast cereals), nuts, herbs and spices, dried soups, dried fruits and vegetables. Pay particular attention to items that have remained in the cupboard for long periods. Foods that are loosely sealed or are in thin wrapping are more likely to be infested than materials in insect resistant containers such as hard plastic or metal.

Indian meal moth may also be found in other materials around the home. Dried dog food and bird seed should be checked. Dried flowers and some craft items that include seeds may be infested. Areas where flour and other materials used in baking may have spilled can support Indian meal moth. Larvae are also known to occur in the stored caches of seeds and nuts that squirrels and other rodents may have around the home.

Infested material should be immediately discarded, used up, or somehow treated to disinfest. Treatments involve using heat or cold to kill any larvae and eggs that may be in the food. Cold treatment requires putting infested items in deep freeze for at least two

or three days. Effectiveness of cold treatment may be improved by alternating freezing treatments with rewarming to room temperatures for a few days. High temperature treatments involve oven heating at around 120 to 140 degrees F for 20 minutes (Somewhat longer intervals are needed if treated items are bulky, requiring longer periods to raise internal temperatures.). Injury to the food is possible with excessively high temperature treatments.

Since insects also can develop on spilled food, thoroughly clean areas where food was stored by vacuuming or sweeping up all spilled food. The thoroughness of the cleaning is important primarily to eliminate food for surviving insects to feed on. The nature of the cleaning agent (soapy water, bleach, etc.) is less important than the permanent elimination of the food.

Heat or cold treated objects are capable of being immediately reinfested as long as Indian meal moths remain in the home so extra care should be taken during this stage. Although adult moths may only live for a week or so, larvae that have recently pupated in hidden areas of the home may also be a potential source of reinfestation. Therefore Indian meal moths must be denied access to all food sources for the length of time that is required to complete the pupal stage plus how long the adult life span can be. A month should be adequate to cover this period.

In the interim, susceptible food items must be stored to prevent reinfestation. Place foods in tightly sealed containers. Food also may be stored in the refrigerator or outdoors until the moths have died out.

Although Indian meal moth can be eliminated, reinfestation are always possible through accidental reintroductions on infested food. Consideration should be given to how bulk foods are stored. Bulk seeds and other commonly infested foods (e.g., bird seed, dog food) should be stored in outbuildings away from the pantry area. Foods in pantries should be stored in containers that are tight-fitting enough to prevent entry by the minute early stage larvae and must be thick enough to prevent it being penetrated by the chewing of late stage larvae.

Use of insecticides within the pantry area is not recommended and will normally give little, if any, additional control in the absence of a thorough clean up and treatment of infested materials. Some household insecticide may allow crack and crevice treatments near food storage areas, but these must be restricted to these site. Never apply insecticides in a manner that allows direct contact with food, food preparation surfaces or food utensils.



Figure 5. Indian meal moths captured at pheromone trap. (Photo by W. Cranshaw.)

There are available traps for Indian meal moth that are baited with an attractant known as a sex pheromone. This is the chemical used by the female Indian meal moth to attract males. Such traps are very useful for identify “hot spots” of infestation. However their ability to control Indian meal moth is highly doubtful, despite occasional claims to this effect by suppliers. This is because the traps only capture males, and usually only a fraction of these. As mated females are not captured, they will continue re-infestation.

Source: Indian Meal Moth by Whitney Cranshaw, Department of Bioagricultural Sciences and Pest Management, Colorado State University.

Current Pests: What Are You Seeing?

Arapahoe, Douglas, & Elbert Counties

Household Insects

Indian meal moth: Adults are most commonly observed flying about homes during early winter.

Fungus gnats: Adults begin to be observed around windows and around the soil of potted plants where they originate.

Boxelder bugs, conifer seed bugs, multicolored Asian lady beetles: Overwintering adults continue to be active in and around homes during warm days.

Fruit flies: Flies from overripe fruit continue to be present in homes.

Denver Metro Area

Household Insects

Indian meal moth: Adults are most commonly observed flying about homes during early winter.

Fungus gnats: Adults begin to be observed around windows and around the soil of potted plants where they originate.

Boxelder bugs, conifer seed bugs, multicolored Asian lady beetles: Overwintering adults continue to be active in and around homes during warm days.

Fruit flies: Flies from overripe fruit continue to be present in homes.

Eastern Plains Counties

Household Insects

Indian meal moth: Adults are most commonly observed flying about homes during early winter.

Fungus gnats: Adults begin to be observed around windows and around the soil of potted plants where they originate.

Boxelder bugs, conifer seed bugs, multicolored Asian lady beetles: Overwintering adults continue to be active in and around homes during warm days.

Fruit flies: Flies from overripe fruit continue to be present in homes.

Credits

Editors: Clyde Wilson, U.S. EPA Region 8; Assefa Gebre-Amlak, CSU Extension; Frank Peairs, CSU Extension; Thia Walker, CEPEP.

Design & layout: Kierra Jewell

“Pest of the month” photo(s): Whitney Cranshaw, CSU Professor & Extension Specialist

Want to subscribe or unsubscribe? Go to: https://lists.colostate.edu/cgi-bin/mailman/listinfo/ccsipm_1

Remember, the CCSIPM listerv is a forum for you to post a message to the entire group! Simply write a message to ccsipm_L@lists.colostate.edu!

Did we miss something? See an error?
Please contact Assefa Gebre-Amlak at:
Assefa.Gebre-Amlak@colostate.edu
(970) 491-2666

El Paso & Teller Counties*Household Insects*

Indian meal moth: Adults are most commonly observed flying about homes during early winter.

Fungus gnats: Adults begin to be observed around windows and around the soil of potted plants where they originate.

Boxelder bugs, conifer seed bugs, multicolored Asian lady beetles: Overwintering adults continue to be active in and around homes during warm days.

Fruit flies: Flies from overripe fruit continue to be present in homes.

High Country Counties*Household Insects*

Indian meal moth: Adults are most commonly observed flying about homes during early winter.

Fungus gnats: Adults begin to be observed around windows and around the soil of potted plants where they originate.

Boxelder bugs, conifer seed bugs, multicolored Asian lady beetles: Overwintering adults continue to be active in and around homes during warm days.

Fruit flies: Flies from overripe fruit continue to be present in homes.

Northern Front Range*Household Insects*

Indian meal moth: Adults are most commonly observed flying about homes during early winter.

Fungus gnats: Adults begin to be observed around windows and around the soil of potted plants where they originate.

Boxelder bugs, conifer seed bugs, multicolored Asian lady beetles: Overwintering adults continue to be active in and around homes during warm days.

Fruit flies: Flies from overripe fruit continue to be present in homes.

Pueblo & Fremont Counties*Household Insects*

Indian meal moth: Adults are most commonly observed flying about homes during early winter.

Fungus gnats: Adults begin to be observed around windows and around the soil of potted plants where they originate.

Boxelder bugs, conifer seed bugs, multicolored Asian lady beetles: Overwintering adults continue to be active in and around homes during warm days.

Fruit flies: Flies from overripe fruit continue to be present in homes.

Southwestern Counties*Household Insects*

Indian meal moth: Adults are most commonly observed flying about homes during early winter.

Fungus gnats: Adults begin to be observed around windows and around the soil of potted plants where they originate.

Boxelder bugs, conifer seed bugs, multicolored Asian lady beetles: Overwintering adults continue to be active in and around homes during warm days.

Fruit flies: Flies from overripe fruit continue to be present in homes.

Tri-River Counties*Household Insects*

Indian meal moth: Adults are most commonly observed flying about homes during early winter.

Fungus gnats: Adults begin to be observed around windows and around the soil of potted plants where they originate.

Boxelder bugs, conifer seed bugs, multicolored Asian lady beetles: Overwintering adults continue to be active in and around homes during warm days.

Fruit flies: Flies from overripe fruit continue to be present in homes.

Source: <http://bspm.agsci.colostate.edu/outreach-button/insect-information/> (Yard/Garden Insect Calendars)